

Sports Injury and Joint Replacement Surgery for the Hip and Knee

ACL, LATERAL COLLATERAL LIGAMENT (LCL), POSTEROLATERAL CORNER RECONSTRUCTION POST-OPERATIVE PROTOCOL

General Guidelines

- Ligament reconstructions are usually performed 1 night in-patient stay
- Supervised physical therapy commences immediately post-operatively. Patients should see their physical therapist as soon as practically possible. Supervised therapy continues for 6-12 months
- Braces are routinely used
- Dr Awwad may alter time frames when indicated.

White compression stockings

The compression stocking helps prevent a blood clot from forming in your legs.

Once you are walking frequently you will no longer need the stocking. If you develop lower leg swelling, tenderness, and/or redness, please contact the Dr Awwad's office or the hospital.

Dressings

The bulky encircling dressings (crepe bandage, velband and pads) may be removed the day after surgery. The small adhesive dressings should be left intact. To shower, bagging the knee in the brace is best. Prior to discharge from hospital, an appointment will be made to see a nurse for a dressing change and wound check between 1-2 weeks post-operatively.

Ice and Elevation

The leg should be intermittently elevated and an ice pack used for 72 hours post-operatively to assist with swelling and pain. Ice packs should be applied for 20-30mins/hr. After 72 hrs, ice packs are no longer required, although can be safely continued and their use is very helpful for pain and swelling.

Pain Medications

The anaesthetist will individualise and organise the appropriate pain relief for patients. Commonly required medication are panadeine forte, tramadol and endone. The routine use of anti-inflammatories is not recommended post-operatively, unless directed by Dr Awwad.

Precautions

Patients should contact Dr Awwad's office or the hospital the operation was performed in, if they develop high temperatures, worsening skin redness, worsening calf, knee or thigh pain and swelling and excessive bleeding or ooze from the incision sites.

Phase I - Maximum Protection (Weeks 0 to 6):

- Brace locked at 0° at all times for six weeks
- Ice and modalities to reduce pain and inflammation
- Use crutches non-weight bearing for 6 weeks.
- Range of Motion
 - Begin aggressive patella mobility
 - ROM 0- 90° out of brace with no varus stress
 - Quadriceps setting, focusing on VMO
 - Multi-plane straight leg raising

Phase II - Progressive Stretching and Early Strengthening (Weeks 6 to 16):

Weeks 6 to 12:

- Continue with modalities to control inflammation
- May begin to progressively increasing weight bearing IN BRACE with set on full ROM (to be worn AT ALL TIMES when weight bearing)
- Range of Motion Full knee extension (no hyperextension)
 - Knee flexion to 120°, progress as tolerated
- Exercises
 - Continue with phase I exercise
 - Bilateral closed kinetic chain squatting
 - Multi-plane open and closed kinetic chain hip strengthening
 - Step-up progression
 - Stationary biking
 - Pool program; focus on ROM
 - Proprioception drills

Weeks 12 to 16:

- Range of Motion
 - Full knee flexion and extension
- Exercises
 - Advance strengthening program progressing to unilateral as tolerated
 - Increase intensity of stationary bike program, may add treadmill walking
 - Advance intensity of pool program; focus on strengthening.

Phase III - Advanced Strengthening and Proprioception Phase (Weeks 16 to 20):

Weeks 16 to 20:

- Range of Motion
 - Full knee flexion and extension with terminal stretch
- Exercises
 - Advance cardiovascular program; no running
 - Increase intensity of closed kinetic chain exercises
 - Advance proprioception drills
 - Initiate gym strengthening progressing from bilateral to unilateral as tolerated
- Leg press, squats, partial lunges, hamstring curls, ab/adduction, calf raises
 - Increase intensity of bike and walking program, may add elliptical trainer
- May begin a pool running program

Phase IV - Advance Strengthening and Plyometric Drill Phase (Weeks 20 to 24):

Weeks 20 to 24:

- Implement a full gym strengthening program; including leg extensions at 30° - 0°
- Begin straight plane running
- Begin controlled lateral functional cord drills

Phase V - Return to Sport and Functional Drills Phase (Weeks 24 to 32):

Weeks 24 to 28:

- Continue with aggressive lower extremity strengthening, cardiovascular training, and flexibility
- Implement multidirectional agility drills
- Begin plyometric drills from bilateral to unilateral as tolerated

Weeks 28 to 32

- Follow-up examination with the physician
- Brace fitting for functional knee brace
- Sports test for return to competition

Do you still have a question about your recovery that has not been answered within this document?

Please contact Dr Awwad's office prior to your surgical date at: awwadadmin@orthosa.com.au

Sometimes we may miss a question that is important to you. If so, please feel free to email us your feedback so that we can improve our service to you and future patients – awwadadmin@orthosa.com.au

Ask **Dr Awwad** to clarify your restrictions prior to surgery to avoid disappointment.



Dr George Awwad

MBBS, FRACS (ORTH), FA ORTH A

Orthopaedic Surgeon

APPOINTMENTS AND ENQUIRIES

P 08 8267 8243 E awwadadmin@orthosa.com.au

W www.drgeorgeawwad.com.au

Ask **Dr Awwad** to clarify your restrictions prior to surgery to avoid disappointment.